

December			
Botanical Name	Common Name	Collection Sites	Collection Tips
<i>Acacia concurrens</i>	CURRACABAH		Pods brown and ready to split.
<i>Acacia nerifolia</i>	WATTLE, OLEANDER		Pods brown and ready to split.
<i>Acacia pendula</i>	MYALL, WEeping (WATTLE)		Pods brown and ready to split.
<i>Acacia salicina</i>	WATTLE, WILLOW		Pods brown and ready to split.
<i>Acacia uncinata</i>	WATTLE, GOLD DUST		Pods brown and ready to split.
<i>Acrotriche aggregata</i>	GROUND-BERRY, RED (a heath)		Red berries
<i>Actinotis helianthii</i>	FLANNEL FLOWER		
<i>Allocasuarina leuhmannii</i>	BULL OAK		
<i>Allocasuarina, littoralis</i>	SHE OAK, BLACK		
<i>Atalaya salicifolia</i>	WHITEWOOD, BRUSH		
<i>Billardiera scandens</i>	APPLE DUMPLINGS	2.Duggan Park, Tba	
<i>Breynia oblongifolia</i>	BREYNIA	1.Charles and Motee Rogers Park, Highfields 3. Duggan Park, Tba	Fruit is ripe when red and soft
<i>Callistemon viminalis</i>	BOTTLEBRUSH, WEEPING RED		
<i>Capparis arborea</i>	CAPER TREE, NATIVE		
<i>Capparis mitchelli</i>	CAPER TREE, NATIVE		
<i>Carex appressa</i>	SEDGE, TALL		
<i>Cayratia acris</i>	WATER VINE, HAIRY		
<i>Chloris truncata C4</i>	WINDMILL, GRASS		Collect when seedhead changes from purple to black, and stems become brittle and straw coloured
<i>Codonocarpus attenuatus</i>	BELLFURIT		

<i>Dianella longifolia</i>	FLAX LILY, SPREADING	2.Duggan Park, Tba	Best results from very fresh seed
<i>Dianella revoluta</i>	ANTHER		Best results from very fresh seed
<i>Dianella nervosa</i>			Best results from very fresh seed
<i>Dichanthium sericeum</i> C4	GRASS, QUEENSLAND BLUE		seeds ripen progressively along seedhead; immediately drop when mature
<i>Diospyros geminata</i>	EBONY, LARGE LEAFED QLD		
<i>Dysoxylum rufum</i>	ROSEWOOD, HAIRY		
<i>Enneapogon nigricans</i> C4	BLACK BOTTLE WASHERS		
<i>Eucalyptus crebra</i>	IRONBARK, NARROW-LEAFED		
<i>Flindersia collina</i>	ASH, LEOPARD		
<i>Geitonoplesium cymosum</i>	LILY, SCRAMBLING		
<i>Glochidion fernandi</i>	CHEESE TREE		
<i>Jacksonia scoparia</i>	DOGWOOD		
<i>Microlaena stipoides</i> C3	WEeping RICE GRASS		
<i>Myoporum montanum</i>	BOOBIALLA, MOUNTAIN		
<i>Rapanea variabilis</i>	MUTTONWOOD		
<i>Psydrax odorata</i> subsp <i>buxifolium</i>	CANTHIUM, BOX-LEAFED	2.Duggan Park, Tba	Collect when black fruit is plump and fresh.
<i>Rhodamnia rubescens</i>	MALLETWOOD, RED		
<i>Rubus rosifolius</i>	RASPBERRY, NATIVE		
<i>Rytidosperma</i> (<i>Austrodanthonia</i>) species C3	WALLABY GRASSES		
<i>Santalum obtusifolium</i>	SANDLEWOOD SHRUB		
<i>Schoenoplectus validus</i>	CLUBRUSH, RIVER		
<i>Senna barclayana</i>	SENNA, PEPPER LEAF		

<i>Sophora fraseri</i>	NECKLACE POD (Listed as vulnerable)		Seed grows best in mother soil. Collect soil at same time as seed.
<i>Sorghum leiocladum</i>	SORGHUM, NATIVE		
<i>Stylidium graminifolium</i>	TRIGGER PLANT ,GRASS		
<i>Stlidium laricifolium</i>	TRIGGER PLANT, TALL		
<i>Themeda triandra C\$</i>	GRASS, KANGAROO	3.Duggan Park, Tba	watch for fertile florets with black twisted awns and solid elmmas; then harvest entire culm and keep in water until seed falls.
<i>Wurmbea dioica</i>	EARLY NANCY (small lily)		Mark site if flowers are found as plants become inconspicuous.